

The nature of assembly bias - III. Observational properties

Ivan Lacerna^{1,2}, Nelson Padilla^{1,3} and Federico Stasyszyn⁴

¹*Departamento de Astronomía y Astrofísica, Pontificia Universidad Católica de Chile, V. Mackenna 4860, Santiago, Chile*

²*Instituto de Astronomía, Universidad Nacional Autónoma de México, A. P. 70-264, 04510, México, D.F., México*

³*Centro de Astro-Ingeniería, Pontificia Universidad Católica de Chile, V. Mackenna 4860, Santiago, Chile*

⁴*Universitäts Sternwarte München, Scheinerstr. 1, D-81679 München, Germany*

ABSTRACT

We analyse catalogues of galaxies from the Sloan Digital Sky Survey (SDSS) and find a weak but significant assembly-type bias, where old galaxies have a higher clustering amplitude at scales $r > 1 h^{-1}\text{Mpc}$ than young galaxies of equal magnitude. When using the definition of age based on the luminosity-weighted stellar age, the difference increases from $-19 < M_r < -18$ to $-22 < M_r < -21$ SDSS galaxies, from a 25 per cent to a 45 per cent effect. In contrast, when using the definition based on the D_n4000 index, the clustering amplitude at large scales shows a weaker signal-to-noise for the assembly bias detection. In semi-analytic galaxies the difference in clustering, using the stellar age and luminosity, is similar at distances beyond $3 h^{-1}\text{Mpc}$, but they overpredict the assembly bias found in SDSS galaxies at smaller scales. We then adapt the model presented by Lacerna & Padilla (Paper I) to redefine the overdensity peak height, that traces the assembly bias such that galaxies in equal density peaks show the same clustering regardless of their age, but this time using observational features. The proxy of peak height for observational data consists in the luminosity inside cylinders of different radii centred in each galaxy. This radius is parameterized as a function of stellar age and magnitude M_r . The best-fitting set of parameters that make the assembly bias practically absent for both simulated and real galaxies are similar ($a = 0.75$, $b = -0.36$ and $a = 1$, $b = -0.36$, respectively). By estimating a weighted average for the values in the range of 1σ for semi-analytic and SDSS galaxies, it is obtained that the set of best-fitting parameters for the radius of the cylinder is $a = 0.76 \pm 0.31$, $b = -0.39 \pm 0.18$. The idea behind the parametrization is not to minimize the bias, but it is to use this method to understand the physical features that produce the assembly bias effect. Even though in this case we consider neighbour galaxies in cylinders in contrast to using a smooth density field, e.g. the dark matter particles distribution, our simulations show that the new method is correlated to the definition of Paper I as both measure the crowding around objects.

Key words: cosmology: large-scale structure of Universe - cosmology: dark matter - galaxies: statistics - galaxies: formation

1 INTRODUCTION

In the current hierarchical clustering models, galaxies form by gas cooling inside virialized cold dark matter haloes in a way that makes their properties depend only on the halo mass in which they form. Recently, results from N -body simulations have shown that the halo clustering at large-scales depends on the halo formation history in addition to the halo mass (Gao et al. 2005; Wechsler et al. 2006; Zhu et al. 2006; Gao & White 2007; Faltenbacher & White 2010). This effect, that is not expected from the excursion set theory, was termed “assembly bias”. For example, Gao et al. (2005) found that low-mass haloes which were assembled at high redshifts are more

strongly correlated than those of the same mass that assembled more recently. Therefore, one could ask whether the assembly bias could also affect galaxy properties.

By means of a semi-analytic model of galaxy formation, Croton et al. (2007) showed that galaxy clustering on large scales is affected by the correlation of halo environment with halo assembly history. Lacerna et al. (2011, hereafter Paper I) found an assembly bias effect in the semi-analytic galaxies from the Lagos, Cora & Padilla (2008) model, where the old population has a higher clustering at large scales than the young population with the same host halo mass.

On the observational side, the existence of the assembly bias is still debated. For example, Skibba et al. (2006) used galaxies from the Sloan Digital Sky Survey (SDSS; York

et al. 2000) to measure marked correlation functions which in addition to the halo model suggest that the correlation between halo formation and environment is not important for the bright galaxy population. Tinker, Wetzel & Conroy (2011) found similar results, indicating that this dependence arises just from the correlation between halo mass function and environment. However, Yang, Mo, & van den Bosch (2006) and Wang et al. (2008) found that groups, selected from the Two Degree Field Galaxy Redshift Survey (2dFGRS; Colless et al. 2001) and SDSS respectively, with red central galaxies are more strongly clustered than groups of the same mass but with blue central galaxies, this effect being much more important for less massive groups. Zapata et al. (2009) found that SDSS galaxy groups of similar mass and different assembly histories show differences in their fraction of red galaxies. Furthermore, Cooper et al. (2010) studied the relationship between the local environment and properties of galaxies in the red sequence. After removing the dependence of the average overdensity on colour and stellar mass by setting a density parameter as a estimator of environment such as it is independent of these galaxy properties, they still found that galaxies with older stellar populations occupy regions of higher overdensities compared to younger galaxies of similar colours or stellar masses. Also, Alonso et al. (2012) show that galaxies in groups in low global densities are prone to show higher star formation compared to galaxies of equal properties and similar host groups but in high density environments. These results show that the concept of assembly bias is applicable to the galaxies in addition to dark matter (DM) haloes and would then influence the physics of galaxy formation. Also, this effect is important when galaxies are used to constrain cosmological parameters. Wu et al. (2008) showed that upcoming photometric catalogues such as the Dark Energy Survey (DES)¹ and the Large Synoptic Survey Telescope (LSST; LSST Science Collaborations et al. 2009)² can infer significantly biased cosmological parameters from the observed clustering amplitude of galaxy clusters if the assembly bias is not taken into account.

Paper I presented a new peak height definition different from the halo virial mass, where semi-analytic galaxies in haloes of a given mass but different age do not show the assembly bias effect in the two-halo regime. This new definition simply adds mass to that of the dark matter halo in a way that makes old, low-mass haloes suffer the most extreme peak height change with respect to what results of using the virial mass as its proxy. In this paper we will test this model using galaxies selected from the SDSS catalogue with the difference that, instead of using the total mass inside spheres of different radii, we will use the total luminosity inside cylinders along the position of each galaxy, which we will use to make a more adequate proxy for a peak height, in a similar way to what was proposed in Paper I. We will discuss the implications of using discrete tracers of the density field that consider neighbour galaxies instead of a smooth density field.

The outline of this paper is as follows. In Section 2 we present the catalogues of galaxies from the SDSS and also

from a semi-analytic model. We characterize the dependence of age on local density in Section 3. We study the assembly bias effect, namely, the large-scale clustering of galaxies of a given luminosity depends on the age, for both the SDSS and semi-analytic galaxies in Section 4. We adopt a modified version of the model from Paper I in Section 5, but using observational features. The nature of the objects that are being considered with the proxy of the overdensity peak height is shown in Section 6, where we discuss the implications of using discrete tracers of the density field that consider neighbour galaxies instead of a smooth density field. The conclusions of our results are presented in Section 7.

2 DATA

2.1 SDSS galaxies

We use galaxies from the spectroscopic SDSS Data Release 4 (DR4, Adelman-McCarthy et al. 2006), which contains $\sim 480,000$ galaxies. After cleaning of repeated galaxies, saturated objects, galaxies close to the edges of the survey, etc., the sample includes $\sim 400,000$ galaxies with measured spectra and photometry in the bands u, g, r, i, z (Fukugita et al. 1996; Smith et al. 2002) that are K -corrected to redshift 0.1 using `kcorrect v4_2` (Blanton & Roweis 2007). The main motivation to use these data is the measured luminosity-weighted ages by Gallazzi et al. (2005), which are essential to this work.³ There are 183,482 galaxies with estimates of stellar age in the magnitude range $-22 < M_r < -18$ (Table 1). We then select four samples of centres and tracers detailed in Tables 1 and 2, respectively. Figure 1 shows the absolute magnitude in the r -band, M_r , as a function of redshift with the choice of volume-limited samples, which avoid observational bias.

The SDSS galaxies will be useful to study whether the assembly bias effect is present in these observed objects thus affecting their properties, as will be shown in Section 4.

2.1.1 Age parameter δ_t

As was the case in Paper I, one of the most important parameters throughout this work is the age. For SDSS galaxies, we use the luminosity-weighted age (or, simply, stellar age) and the D_n4000 index (Balogh et al. 1999). The latter is defined as

$$D_n4000 = \frac{\overline{F}_\nu(4000 - 4100 \text{ \AA})}{\overline{F}_\nu(3850 - 3950 \text{ \AA})}, \quad (1)$$

where $\overline{F}_\nu$ is the average flux density in the specified bands (in the rest frame), and it is smaller for young stellar populations than for old, metal-rich galaxies. In our samples there are 183,482 galaxies with estimates of stellar age (Table 1) and 205,076 with D_n4000 , both in the magnitude range $-22 < M_r < -18$.

³ Luminosity-weighted ages are obtained from <http://www.mpa-garching.mpg.de/SDSS/DR4/Data/stellarmet.html> which includes the DR4 version of the tables used in Gallazzi et al. (2005).

¹ See <http://www.darkenergysurvey.org/>.

² See <http://www.lsst.org/lsst/>.

Table 1. Centre samples used to measure the cross-correlation functions. The second column indicates the number of objects with luminosity-weighted ages, the third column corresponds to the magnitude bin, and the last column indicates the redshift range of each volume-limited sample (see Fig. 1).

	N	M_r	redshift
centres sample 1	10 953	$-19 < M_r < -18$	$0.010 < z \leq 0.050$
centres sample 2	41 167	$-20 < M_r < -19$	$0.012 < z \leq 0.078$
centres sample 3	78 632	$-21 < M_r < -20$	$0.020 < z \leq 0.120$
centres sample 4	52 730	$-22 < M_r < -21$	$0.035 < z \leq 0.175$

Table 2. Same as Table 1 but for tracers. In this case, the third column corresponds to a magnitude limit.

	N	M_r	redshift
tracers sample 1	33 907	$M_r < -18$	$0.010 < z \leq 0.050$
tracers sample 2	77 114	$M_r < -19$	$0.012 < z \leq 0.078$
tracers sample 3	103 135	$M_r < -20$	$0.020 < z \leq 0.120$
tracers sample 4	60 180	$M_r < -21$	$0.035 < z \leq 0.175$




Figure 1. Absolute magnitude M_r as a function of redshift for SDSS galaxies. Four volume-limited samples of centres and tracers are chosen to perform cross-correlation functions without observational bias. Objects inside the rectangles in red, blue, green, and cyan correspond to the centre samples 1, 2, 3, and 4 in Table 1. For each sample, the tracers (see Table 2) are in the same range of redshift (delimited by the vertical lines) as centres but also include brighter objects.

To study the assembly bias in galaxies, where old objects may have a different clustering than young objects of the same mass or luminosity, it is not convenient to work directly with the stellar age, for example, because this quantity correlates with the mass or the luminosity of galaxies. For instance, luminous galaxies have, on average, older stellar




Figure 2. Stellar age as a function of the absolute magnitude M_r for SDSS galaxies, which correspond to the volume-limited samples described in Table 1. Due to the large number of available galaxies, only 10,000 of them, randomly chosen, are plotted as black points. Red squares show the medians for each magnitude bin. Error bars correspond to the 10 and 90 percentiles of the stellar age distribution. The median population of fainter galaxies is younger than that of the brighter objects.

populations (see Figure 2) as indicated by their luminosity-weighted age. In Paper I, we presented a definition of age which is independent of the mass or the luminosity through the δ_t dimensionless parameter

$$\delta_{t_i} = \frac{t_i - \langle t(M_r) \rangle}{\sigma_t(M_r)}, \quad (2)$$

where for the i th galaxy, t_i is its stellar age, $\langle t(M_r) \rangle$ is the median stellar age as a function of M_r , and $\sigma_t(M_r)$ is the dispersion around the median obtained from the 10 and 90 percentiles of the distribution (error bars in Fig. 2). Then, objects with positive (negative) values of δ_t correspond to old (young) populations with respect to the median distribution of stellar ages of their luminosity. The scatter in the observational stellar age can be higher than the stellar age obtained in simulated galaxies such as the model described in Section 2.2.

We also use D_n4000 in Equation (2) as a proxy for age considering its dependence on the absolute magnitude M_r (similar to Figure 2).

2.2 The numerical simulation

We use synthetic galaxies from the SAG2 model by Lagos, Cora & Padilla (2008, see also Lagos, Padilla & Cora 2009), who combine a cosmological N -body simulation of the concordance Λ CDM universe and a semi-analytic model of galaxy formation to follow the evolution of the baryonic component of haloes accross cosmic history. The numerical simulation consists in a periodic box of $60 h^{-1}$ Mpc on a side that contains 256^3 dark matter particles with a mass resolution of $\sim 10^9 h^{-1} M_\odot$. The cosmology used is $\Omega_{tot} = 1$, $\Omega_m = 0.28$ (with a baryon fraction of 0.16), $\Omega_\Lambda = 0.72$, $\sigma_8 = 0.9$, $h = 0.72$, and spectral index $n_s = 1$. Dark matter haloes are identified as self-bound structures that contain at least 10 particles using a friends-of-friends (FOF) algorithm (Davis et al. 1985). Another algorithm (SUBFIND; Springel et al. 2001) is applied to these groups in order to find substructures with at least 10 particles. Typically, the most massive subhalo is considered the main subhalo, while the other subhaloes are satellites within a FOF halo. The galaxy population in the semi-analytic model is generated using the merger histories of dark matter haloes and their embedded subhaloes. The virial mass for galaxies corresponds to the virial mass of their respective DM substructures.

For the simulated galaxies, we will use the mass-weighted stellar age in equation (2). Magnitudes in the SAG2 semi-analytic model are in the Johnson system. Therefore, we apply a correction to transform these magnitudes to the *ugriz* system to compare with SDSS galaxies using the equations of Jester et al. (2005) where, for objects with $R-I < 1.15$ and $U-B < 0$,

$$r = V - 0.46(B - V) + 0.11 \quad (3)$$

and, for objects with just $R-I < 1.15$,

$$r = V - 0.42(B - V) + 0.11. \quad (4)$$

Figure 3 shows scatter plots of halo DM mass (virial mass as explained above and friends-of-friends in the left-hand and right-hand panels, respectively) vs. M_r . Both definitions

Table 3. Mean virial mass M_{vir} in units of $h^{-1} M_\odot$ for old ($\delta_t > 0.05$) and young ($\delta_t < -0.05$) galaxies of a given magnitude M_r using the SAG2 semi-analytic model (third column) and the semi-analytic model of De Lucia & Blaizot (2007) (fourth column). Given uncertainties the differences are not significant.

M_r	Ages	$\log\langle M_{vir} \rangle^a$	$\log\langle M_{vir} \rangle^b$
$-19 < M_r < -18$	Old	10.6	10.7
	Young	10.8	10.8
$-20 < M_r < -19$	Old	11.1	11.0
	Young	11.1	11.1
$-21 < M_r < -20$	Old	11.6	11.5
	Young	11.5	11.5
$-22 < M_r < -21$	Old	12.2	12.2
	Young	12.0	11.9

^aSAG2.

^bDe Lucia & Blaizot (2007).

show a strong correlation with the inferred r -band absolute magnitude, which is strong for the virial mass, thus we are able to use M_r as a proxy of mass, specially that of the substructure (Conroy et al. 2006; Simha et al. 2012).

One may think that old galaxies have larger mass-to-light ratios than young galaxies of equal luminosity. Table 3 shows the mean virial mass for old and young galaxies of equal luminosities using the SAG2 semi-analytic model and the semi-analytic model of De Lucia & Blaizot (2007).⁴ Therefore, there is no a strong dependence regarding the virial mass when splitting the old and young galaxies for a given luminosity. Furthermore, in only one out of the four samples of our model, the halo mass is lower for the older population, which goes in the opposite direction to the common sense interpretation of the cuts in age. This is due to the fact that we are looking at narrow ranges in luminosity.

Figure 4 shows that the scatter between virial mass and the stellar mass is similar to that using M_r . Therefore, there is little difference in using luminosity or stellar mass as a proxy for mass (at least that of the substructure).

3 DEPENDENCE OF RELATIVE AGE ON LOCAL DENSITY

To adapt the procedure of Paper I to galaxies, we need to understand the age-density aspect of the assembly bias in both simulations and observations. Figure 5 shows a relative stellar age map in our simulation, where the relative age δ_t is calculated with respect to virial mass using equation (2) of Paper I. The colours represent different average values of δ_t in grids conformed by cells of $2 h^{-1}$ Mpc on a side in a slice $5 h^{-1}$ Mpc thick of our simulation. In general, regardless if the relative stellar age is calculated with respect to virial mass

⁴ This model made use of the Millennium Simulation (Springel et al. 2005). We find that this quantity is very similar for galaxies of different age for a given magnitude using both models.




Figure 3. Scatter plots of mass versus the inferred absolute magnitude M_r for galaxies in the simulation. Left: as a function of virial mass, M_{vir} . There is a strong correlation between M_{vir} and M_r , where more massive objects are brighter. Right: the mass is given by the friends-of-friends (FOF) approach, which is defined as the total mass of a FOF halo (i.e. a satellite galaxy has a M_{fof} equal to the the central galaxy). This latter definition blurs the correlation seen in the left panel.




Figure 4. Scatter plots of virial mass versus the stellar mass for galaxies in the numerical simulation. As well as the luminosity, there is a strong correlation between M_{vir} and stellar mass.

o luminosity, values of $\delta_t \geq 0.05$ correspond to relatively old populations, while $\delta_t \leq -0.05$ to young populations. As can be seen, the latter appear less clustered than the old population even after, in this case, the mass dependence is removed using the definition of δ_t .

We quantify this behaviour using the D_n4000 index. The distribution of D_n4000 for our SDSS galaxies is shown in the left-hand panel of Figure 6 for different ranges of ab-




Figure 5. Age maps using the δ_t parameter based on the stellar age with respect to virial mass in a slice of the numerical simulation. The colour bar indicates the different ranges in δ_t .

solute magnitude. We see a clear bimodality for all of the magnitude bins plotted except for the brightest one, probably because most of these objects are part of the ‘red sequence’ (poor gas galaxies with low star formation rate). The vertical-dotted line shows the cut that divides the two populations; active galaxies ($D_n4000 < 1.65$) and quenched galaxies ($D_n4000 > 1.65$). Note that it is very similar to the

condition used by Tinker, Wetzel & Conroy (2011), where quenched objects have $D_n4000 > 1.6$. The right-panel of Fig. 6 shows the fraction of quenched SDSS galaxies as a function of the large-scale environment. The latter is defined by counting the number of neighbour galaxies inside a sphere of $10 h^{-1}$ Mpc radius around each galaxy (see Muldrew et al. 2012 for a revision of different environment definitions including this one). We also restrict to neighbours above the magnitude limit for each sample. We see that the fraction of quenched objects increases to higher densities. Our results are qualitatively similar to those found by Tinker, Wetzel & Conroy (2011) using this index.

The trend that old objects are located in regions of higher densities than young objects of equal mass or magnitude is also readily seen in the two-point correlation function (see Section 4, for example), where the ones with the higher clustering at large scales are those located on higher density regions, which could be indicating the locations of higher initial density peaks. The correlation function, $\xi(r)$, has the advantage that it obtains an accurate characterization of the spatial distribution of objects from kpc scales out to Mpc scales by simply measuring the excess of pairs at a distance r with respect to a random distribution. Therefore, $\xi(r)$ is a useful tool to study the large-scale environment. The overall results will support that older populations are located in higher densities compared to younger ones regardless the luminosity.

High density environments play a key role in properties such as the formation age. Later in this work we will use this density-age connection to obtain a peak-height proxy. We will show that the crowding around objects helps us trace the assembly bias effect as well as the smooth density field approach introduced in Paper I.

4 ASSEMBLY BIAS IN SDSS GALAXIES

In this section, we will test whether SDSS galaxies show an assembly type bias by using the two-point correlation function for the centres and tracers in the samples of Tables 1 and 2. Due to the redshift space distortions, we have to calculate first the cross-correlation in two coordinates $\xi(\sigma, \pi)$, where σ is perpendicular and π is parallel to the line of sight. Figure 7 shows the resulting $\xi(\sigma, \pi)$ for the old and young galaxies in the magnitude range $-21 < M_r < -20$. As can be seen there is a subtle difference between both populations due to their slightly offset clustering amplitudes (c.f. Figure 8), but also possibly due to different finger-of-god and infall effects. Given that the differences in infall are too subtle, we will not use these profiles as estimators of the assembly bias in contrast to Paper I.⁵

I use the jackknife technique to estimate the variance error in the calculation of $\xi(\sigma, \pi)$. In particular the original data is divided in 20 subsamples. The correlation function is calculated by discarding one of the subsamples at a time, and then it is compared with $\xi(\sigma, \pi)$ obtained for the whole sample in order to build the error estimator.

We integrate over the π component to obtain the projected correlation function $\omega(\sigma)$,

$$\omega(\sigma) = 2 \int_{\pi_{min}}^{\pi_{max}} \xi(\sigma, \pi) d\pi, \quad (5)$$

where $\pi_{min} = 0.1 h^{-1}$ Mpc and $\pi_{max} = 30 h^{-1}$ Mpc. This can be inverted to obtain the real-space cross-correlation function using the relation described by Saunders, Rowan-Robinson & Lawrence (1992, see also Paz, Staszczyn & Padilla 2008),

$$\xi(r) = \frac{-1}{\pi} \sum_{j \geq i} \frac{\omega(\sigma_{j+1}) - \omega(\sigma_j)}{\sigma_{j+1} - \sigma_j} \ln \left(\frac{\sigma_{j+1} + \sqrt{\sigma_{j+1}^2 - \sigma_i^2}}{\sigma_j + \sqrt{\sigma_j^2 - \sigma_i^2}} \right). \quad (6)$$

The panels of Figure 8 show $\xi(r)$ for old and young objects, selected using ranges in δ_t , for samples of different luminosities. Error bars are calculated taking into account the error estimate for $\xi(\sigma, \pi)$ and using equations (5) and (6). We systematically find that the old galaxies show a higher clustering relative to the young population at large scales ($r > 1 h^{-1}$ Mpc) for different ranges in M_r . This difference typically increases toward brighter objects from a 25 per cent to a 45 per cent effect. Therefore, SDSS galaxies of equal luminosity show a significant dependence of clustering amplitude with the age since the differences are over a 20 per cent effect. As can be seen from Table 3, the mass-to-light ratio M/L is not higher for old galaxies at fixed luminosity in the simulation, so that this dependence of clustering amplitude with age is the assembly bias effect.

This result is in agreement with Wang et al. (2008), Zapata et al. (2009) and Cooper et al. (2010) who find an assembly-type bias in galaxies. Other works, however, mention that the environmental dependence in galaxies is not important at large-scales. For example, Tinker, Wetzel & Conroy (2011) found that the fraction of quenched galaxies increases with the large-scale environment, but this arises just from the dependence of the halo mass on the environment. They use the D_n4000 parameter to classify a quenched galaxy ($D_n4000 > 1.6$). We find that when using this index to separate active/young and quenched/old objects, the difference in clustering amplitude at large scales due to assembly bias is not clear. The left-panel of Figure 9 shows the ratio of the correlation function of old galaxies to the total sample of a given magnitude and the same ratio but using young galaxies, where the determination of age is based on D_n4000 in Equation (2). The difference in the clustering amplitude between the old and young population is very noisy using this estimator for the age, in contrast to the results obtained using the luminosity-weighted stellar age shown in the right panel (see also lower boxes of Fig. 8). Therefore, the D_n4000 index is less adequate to show clearly the environmental dependence at large scales, even when we have 12 per cent more galaxies with this index than with the stellar age (c.f. Section 2.1). Probably, this is due to the fact that the D_n4000 index is related with star formation by re-

⁵ Except in Section 6 where we reproduce the method used in Lacerna & Padilla (2011).




Figure 6. Left: distribution of the D_n4000 index for SDSS galaxies, normalized by 10,000 of them, for four bins in magnitude. We see a clear bimodality for all except the brightest one. The vertical-dotted line is our cut to define quenched galaxies, $D_n4000 > 1.65$. Right: fraction of quenched SDSS galaxies as a function of the environment for different magnitude bins.




Figure 7. Cross-correlation functions in the directions parallel and perpendicular to the line of sight for old (left panel) and young (right panel) SDSS galaxies in the magnitude range $-21 < M_r < -20$. The colour in the figure key indicates the amplitude of the correlation function. The three contour levels, from large to small separations, correspond to $\xi(\sigma, \pi) = 0.5, 1.0$ and 5.0 , respectively.

cent bursts in contrast to the luminosity-weighted age which covers an important range of star formation histories.⁶

4.1 Cross-correlation functions for the simulated galaxies

We use the $z = 0$ model galaxies and then select four samples of centres and tracers in the same ranges of M_r indicated in Tables 1 and 2, respectively. Figure 10 shows the results of the different cross-correlation functions. Error bars are calculated using the jackknife method, which has been shown as a robust error estimator for cosmological samples. In general, this method consists in dividing the sample in parts.

⁶ Gallazzi et al. (2005) includes the D_n4000 index among other spectral features to constrain the luminosity-weighted age.




Figure 8. Cross-correlation functions in real space for SDSS galaxies (see Section 4 for details). The bins in magnitude M_r are indicated in each panel. Old and young galaxies are shown in red circles and blue squares, respectively. Error bars were calculated using the jackknife method. The lower box in each panel corresponds to the ratio between the correlation function of the old population and the total population of the selected sample and, also, between the young population and the total one (dot-dashed red and dotted blue lines, respectively). The error of the ratio between the $\xi(r)$ of the old and young objects is shown as a shaded region around the unit ratio.

The correlation function is calculated by discarding one of the subsamples at a time, $\xi_i(r)$, and then it is compared with the correlation function obtained for the whole sample $\xi(r)$ as follows

$$\Delta^2 = \frac{1}{N} \sum_{i=1}^N [\xi_i(r) - \xi(r)]^2, \quad (7)$$

where N is the total number of subsamples. In particular, samples selected from the numerical simulation are divided in ten parts, $N = 10$ (for samples selected from SDSS data,

$N = 20$). As can be seen from these panels, old galaxies (red circles) are more strongly clustered than the young objects (blue squares), with average differences of up to a factor of two in the clustering amplitude at large scales, $r > 1 h^{-1}$ Mpc.

In order to compare the results from real and simulated galaxies, Figure 11 shows the ratio $\xi(r)$ of the old galaxies to that of the young galaxies from the simulation and SDSS-DR4 (solid and dash lines, respectively). Both ratios are similar for distances beyond $3 h^{-1}$ Mpc, especially for brighter




Figure 9. Ratios between the correlation function of the old SDSS galaxies and the total population of each selected sample (dot-dashed red lines) and, also, between the young SDSS galaxies and the total one (dotted blue lines), where the definition of age for the galaxies is based on the D_n4000 index (left panel) and the luminosity-weighted stellar age (right panel). Errors are not shown for clarity.

magnitudes. The clustering of old objects is typically thirty per cent higher than that of young galaxies. At lower scales the ratio is higher for the semi-analytic galaxies, while it remains constant for the SDSS galaxies. This difference may be explained, for instance, by the simplified method of circular orbits in the simulation that could affect the profiles on the one-halo term (Lagos, Cora & Padilla 2008). Note that Guo et al. (2011) found that the projected autocorrelation functions for their semi-analytic galaxies are higher than for observations at scales $r < 1$ Mpc. They suggest what this is due to the fact that their assumed fluctuation amplitude, $\sigma_8 = 0.9$, the same value used in this work, is too high. The latest results of WMAP7 (Komatsu et al. 2011) estimate $\sigma_8 = 0.809 \pm 0.024$. Furthermore, Figure 11 shows the ratios of two correlation functions, pointing to a stronger overprediction for old galaxies which reside in denser regions, which is consistent with Guo et al. (2011), who also find that the simulation overpredicts the autocorrelation functions for passive galaxies, while underpredicts it for active galaxies.

5 OVERDENSITY PEAK HEIGHT PROXY USING GALAXY LUMINOSITIES

In Paper I, we presented an approach to trace the assembly bias effect. This consisted in measuring the total mass inside spheres of different radii around semi-analytic galaxies, which in some cases could be larger than the virial mass of the host dark matter halo. Using two free parameters to introduce a dependence of this radius on mass and age, this model defined a new overdensity peak height for which the large-scale clustering of objects of a given mass did not depend on the age. We then analyzed the changes of this peak height proxy, compared to virial masses. We typically found that low-mass, old galaxies have larger peak heights than

young galaxies of equal virial mass, which implies that the environment is more dense for old objects. They are preferentially surrounded by massive neighbour haloes, which is in agreement with other works that mention old, low-mass haloes may suffer a truncated growth due to massive neighbour haloes (Wang, Mo & Jing 2007; Dalal et al. 2008; Hahn et al. 2009). Therefore, the parametrization of the radius for the sphere to measure the environment is a useful tool to understand what is behind the assembly bias.

However, it is difficult to obtain the underlying mass information for real galaxies. For this reason, the aim of this section is to design a similar procedure, but using the luminosity of galaxies. We take advantage of the fact that galaxies of equal virial mass have similar luminosities (c.f. Section 2.2). The idea behind our work is not to minimize the bias, but it is to use this method to understand the physical features that produce the assembly bias.

Instead of using spheres as in Paper I, we measure the total luminosity (which correlates with total substructure mass) inside a cylinder centred on each galaxy along the line-of-sight to take into account the limitations imposed by the redshift space distortions. The radius of the cylinder for each galaxy in h^{-1} Mpc units is parameterized as

$$r = a \delta_t + b (M_* - M_r), \quad (8)$$

where the characteristic magnitude is $M_* = -20.5$ (Zehavi et al. 2005). The free parameters are a and b . The length of the cylinder is $\Delta v = \pm 500$ km/s centred in each galaxy.⁷ We add the luminosity L of these objects to ob-

⁷ The choice of cylinder length ± 500 km/s corresponds to a total length of $5 h^{-1}$ Mpc in the z -coordinate of the simulation.




Figure 10. Same as Figure 8 but for the galaxies in the simulation. Error bars were calculated using the jackknife method (see equation 7) by dividing the sample in ten parts. The error of the ratio between the $\xi(r)$ of the old and young objects is shown as a shaded region around the unit ratio in each lower box.

tain the total luminosity in this volume using the relation $M_r - M_* = -2.5 \log(L/L_*)$. When the radius is negative the new magnitude of the central object remains unchanged, i.e. $M'_r = M_r$. If this radius is greater than 3 Mpc, the object will not be considered for further analysis (for the best-fitting set of parameters, none of the galaxies exceed this limit in radius).

Notice that this cylinder will include neighbour galaxies and therefore this will be related to the crowding of the environment of the galaxy, a measure of the local density, but with an adaptive volume size depending on the galaxy age and luminosity. This procedure to trace the assembly bias will use correlation functions of relatively old and young

galaxies to fit the parameters of the cylinder radius, but this parametrization will only be said to correlate with the correction of Paper I (missing peak mass) if the excess of crowding in galaxies correlates with the excess of mass found in haloes reported in our previous paper. In order to demonstrate that this is the case, we first need to find the parameters that trace the assembly bias using this measurement of crowding, and then to check whether the relation between crowding and missing peak mass is present. In contrast to Paper I, we do not use the infall velocity profile because the velocity inferred from redshift space distortions is noisy compared to the velocity from numerical simulations.




Figure 11. Ratio between the correlation functions of the old population and that of the young population for the different ranges in luminosity indicated in each panel. Dashed lines (open squares) represent the real-space results of the SDSS galaxies as shown in Fig. 8. Solid lines (filled circles) correspond to those from the simulated galaxies (see Fig. 10). Dotted horizontal lines show the unit ratio (no assembly bias).

5.1 Simulated galaxies

Our aim is to affect the model with the observational problems for which we define the z -axis of the simulation as the line-of-sight and turn the z -coordinate into a recession velocity, $v'_z = 100z + v_z$, where v_z is the velocity component in the z -coordinate. When measuring the luminosity in each cylinder in the numerical simulation, we include all the objects with stellar masses higher than $10^9 h^{-1} M_\odot$ and magnitude M_r brighter than the limit given for each sample in Table 2.

We calculate the cross-correlation functions between samples of a given new magnitude M'_r and tracer galaxies that satisfy $M_r < -18$. We also select subsamples of old and young objects using the proxy for the age, δ_t , as measured in Section 2.1.1 but using M'_r .

In order to find the best-fitting parameters that do not show an age dependence of the clustering, the reduced χ^2 for the correlation function statistics is defined as

$$\chi^2_{\xi(r)} = \frac{1}{4} \sum_i \left(\frac{1}{n_{dof}} \sum_r \frac{[N_{young}(r) - N_{old}(r)]^2}{\Delta_{N(r)}^2} \right), \quad (9)$$

and is calculated within the range $0.8 \leq r/h^{-1} \text{ Mpc} \leq 10$, mostly in the two-halo term; N_{young} is the number of neighbours for young galaxies, whereas N_{old} is the same quantity for old objects. The number of neighbours is defined as $N(r) = \langle N_t(r) \rangle \xi(r) + \langle N_t(r) \rangle$, where $\langle N_t(r) \rangle$ is the mean number of tracers (see Paper I for details). The error is $\Delta_{N(r)}^2 = \Delta_{N_{young}}^2 + \Delta_{N_{old}}^2$, calculated as the relative error of the number of neighbours. The symbol n_{dof} denotes




Figure 12. Solid curves show the ratio between the correlation functions of the old and young populations of semi-analytic galaxies for different ranges in magnitude M'_r indicated in each panel for the best-fitting values $a = 0.75$ and $b = -0.36$. Dotted horizontal lines show the unit ratio. For comparison, solid circles (dashed lines) correspond to the ratio shown in Fig. 11 for the simulated galaxies using M_r . Note that this comparison is just for reference because the adopted magnitudes are different.

the number of degrees of freedom. The value $\chi^2_{\xi(r)}$ is the average over four magnitude bins selected between the 10 and 70 percentiles of the M'_r distribution. This particular definition of goodness of fit avoids selecting parameters favoured by large uncertainties that can induce spuriously good fits.

The best-fitting parameters corresponding to the highest likelihood in the full parameter space are $a = 0.75$ and $b = -0.36$ ($\chi^2 \sim 0.51$). Figure 12 shows the ratios of the correlation functions of old and young objects using these values for different magnitude ranges. The assembly bias is practically absent in the numerical simulation using this formalism (compare with solid lines in Figure 11). The average differences in clustering amplitude are smaller than a 10 per cent for each range in M'_r .

Note that we do not include objects with fainter magnitudes than the limit given by the central galaxy in the cylinder. Figure 13 shows the true total magnitude if we also include these neighbour galaxies vs. M'_r . There is a strong relation between both magnitudes with a small scatter that is more important for fainter objects. Furthermore, if we use all the neighbour galaxies, ignoring the restriction of the flux limited catalogues, the best-fitting parameters are $a = 0.5$ and $b = -0.36$ ($\chi^2 \sim 1.01$); the parameter a changes slightly to a lower value, while the parameter b is the same. Figure 14 shows the likelihood functions for the no-flux limit approach (dashed lines) and those of the magnitude M'_r (solid lines). Notice that the maximum likelihood in this figure occurs for parameter values that do not necessarily coincide with those of the maximum likelihood in the full parameter




Figure 13. Magnitude M'_r that accounts for the biased luminosities around central simulated galaxies in cylinders of radius $r = a\delta_t + b(M_* - M_r)$, with free parameters $a = 0.75$ and $b = -0.36$ (see Section 5.1 for details), as a function of the true total magnitude in the r -band if the contribution of all the semi-analytic galaxies located in these cylinders is included. We find a strong relation between both magnitudes with a small scatter that is more important for fainter objects. These central semi-analytic galaxies are in the range $-22 < M_r < -18$.

space. We find that both approaches for the luminosity in the r -band obtain similar maximum likelihood values consistent within a 1σ confidence (indicated by the intersections of the parameter distributions with the dotted line). This justifies our use of M'_r as a good estimator of the total luminosity around galaxies in the range $-22 < M_r < -18$ with the best-fitting values $a = 0.75$ and $b = -0.36$, avoiding the complications introduced by the flux limit nature of the observational catalogs.

In the next section we show a similar approach to redefine the overdensity peak height using luminosities applied to SDSS galaxies.

5.2 SDSS galaxies

Figure 15 illustrates the volume-limited samples used in this section in order to avoid observational bias, where M'_r is the redefined luminosity. We have hundreds of volume-limited samples such as Figure 15 given by the different combinations of the parameters a and b , thus this was done automatically in order to optimize the number of objects in selected subsamples in luminosity. The procedure to define these subsamples is as follows. For each cut in M'_r , we count the number of galaxies in four ranges of redshift indicated in the last column of Table 1. We then select the range that is most populated that defines faint and bright limits in M'_r and a maximum redshift, z_{lim} . All the galaxies located between the magnitude limits of this range with $z \leq z_{lim}$ are




Figure 14. Marginalized likelihood functions for the free parameters a and b (top and bottom panels, respectively) using the magnitudes of semi-analytic galaxies described in Fig. 13 and Section 5.1, the true total magnitude in the r -band (dashed lines) and M'_r (solid lines). Both approaches for the luminosity of cylinders of radius r (Equation 8) obtain similar maximum values in the range of 1σ (intersection of distributions with the horizontal dotted line). We also include the results for the magnitude M'_r for SDSS galaxies described in Section 5.2 (dot-dashed lines).

selected as members of a centre sample, and all the galaxies brighter than the faint-end limit of this range with $z \leq z_{lim}$ are selected as a tracer sample in order to perform cross-correlation functions. For example, the samples shown in Figure 15 correspond to the parameters $a = 0.5$, $b = -0.36$. In this case, for the first cut in magnitude, we select the galaxies with $-19.92 \leq M'_r \leq -19$ and $z \leq 0.078$ (red dots) as the first centre sample. Its respective tracer sample are the galaxies that satisfy $M'_r \leq -19$ and $z \leq 0.078$. For the last cut in magnitude, we select the centre sample (cyan dots) using the conditions $-23.39 \leq M'_r \leq -22.23$ and $z \leq 0.078$. Galaxies for the respective tracer sample satisfy $M'_r \leq -22.23$ and $z \leq 0.078$.

We again select subsamples in δ_t using the distribution of the luminosity-weighted stellar age as a function of M'_r . The correlation functions are obtained in redshift space as




Figure 15. Magnitude M'_r of SDSS galaxies as a function of redshift for the set of parameters $a = 0.5$, $b = -0.36$. Four volume-limited samples of centres and tracers are chosen to perform cross-correlation functions without observational bias. Further details of the selection of centres (in colour regions) and tracers in Section 5.2.

shown in Figure 16. We estimate their real space counterparts as described in Section 4.

The number of neighbour objects in equation (9) using SDSS samples is at least an order of magnitude lower than that in the simulation, so that the statistic to perform the reduced χ^2 is affected by significant noise. The number of neighbours is typically smaller than dozens of SDSS objects, whereas this number is greater than a hundred for simulated objects. This is due to the way the number of neighbours is calculated in equation (9), which is proportional to the mean number of tracers at a distance r , $\langle N_t(r) \rangle \propto V(r)/V_{tot}$, where the total volume of the simulation is $V_{tot} \sim 10^5 \text{ Mpc}^3$ and, for example, the total volume covered by the SDSS sample at $z = 0.175$ is $V_{tot} \sim 10^7 \text{ Mpc}^3$. For this reason, the reduced χ^2 in this case (M'_r for SDSS galaxies) is defined as

$$\chi^2_{\xi(r)} = \frac{1}{4} \sum_i^4 \left(\frac{1}{n_{dof}} \sum_r \frac{[\xi_{old}(r) - \xi_{young}(r)]^2}{\Delta^2} \right)_i, \quad (10)$$

where $\xi_{old}(r)$ and $\xi_{young}(r)$ is the result for old and young SDSS galaxies, respectively. The error is $\Delta^2 = \Delta_{old}^2 + \Delta_{young}^2$, where the first term is the jackknife error for $\xi_{old}(r)$ and the second for $\xi_{young}(r)$. This statistics is estimated within the range $0.7 \leq r/h^{-1} \text{ Mpc} \leq 10$ and averaged over the four magnitude bins selected in M'_r .

For the SDSS galaxies, the best-fitting values are $a = 1$, $b = -0.36$. The marginalized probability distributions for a and b are shown as dot-dashed lines in Figure 14, and as can be seen there is a good agreement with the distributions obtained from our simulation. It is remarkable that the parameter a has a value only slightly greater compared to that obtained for the galaxies in the simulation ($a = 0.75$), which lies within the uncertainties of this parameter, and the value of b is equal in both cases. The correlation func-

tion ratios using these values are shown in Figure 17. Again, the assembly bias is almost not present using the proxy for the overdensity peak height at $r > 1 h^{-1} \text{ Mpc}$ (average differences in clustering amplitude are smaller than a 10 per cent for each range in magnitude M'_r), thus confirming that our approach is able to define an initial peak-height proxy using the crowding of the region where galaxies lie, in a similar manner to that implemented in Paper I, and that SDSS galaxies show a similar behaviour to ΛCDM semi-analytic galaxies since their best-fitting sets of parameters are comparable.

In this context, as mentioned above, Figure 14 shows the marginalized likelihood distributions for the free parameters a and b of equation (8) for three cases: the new magnitude M'_r for simulated galaxies, the true total magnitude in the r -band using also simulated galaxies and M'_r for SDSS galaxies. By estimating a weighted average for the values in the range of 1σ for the three distributions, it is obtained

$$a = 0.76 \pm 0.31$$

$$b = -0.39 \pm 0.18.$$

The best estimates for the a parameter are always positive, whereas those for the b parameter are negative when defining an overdensity peak height proxy using luminosities. The next section will discuss the properties of this proxy of peaks when using galaxy luminosities.

6 PEAKS DEFINED USING LUMINOSITY

The redefinition of the overdensity peak height using galaxy luminosities is able to trace the assembly bias (Section 5). Table 4 shows the median radius of equation (8) in physical units of $h^{-1} \text{ kpc}$ for the four different bins in M_r using the simulated galaxies. The results are shown for two sets of best-fitting values, the M'_r case ($a = 0.75$, $b = -0.36$) and the true total magnitude in the r -band ($a = 0.5$, $b = -0.36$). The radius of the cylinder for all the objects decreases to higher luminosities with very similar sizes for both sets of parameters. This is in agreement to what was discussed in Paper I, where the radius decreases to more massive objects, due to the correlation between luminosity and virial mass. Bear in mind that the radius can be negative, $r < 0$, since both the first and second term of equation (8) can be negative, which implies $M'_r = M_r$. The results are also split according to the classification of old and young populations after performing the redefinition of luminosity for these simulated galaxies. As well as Paper I, old objects have higher radii than young objects (of equal luminosity in this case), thus suggesting stronger environmental dependences for old populations. The median sizes for both sets of parameters are similar for young galaxies, with differences smaller than $37 h^{-1} \text{ kpc}$, while these differences for old galaxies range from $8 h^{-1} \text{ kpc}$ out to $200 h^{-1} \text{ kpc}$ toward brighter magnitudes.

Table 5 shows the median radius of equation (8) using the SDSS galaxies. In general, there is a good agreement of the radius sizes with respect to the M'_r case (see Table 4), with differences among them smaller than $22 h^{-1} \text{ kpc}$ for




Figure 16. Same as Figure 7 also for SDSS galaxies, but using the best-fitting values $a = 1$, $b = -0.36$ in equation (8) for galaxies in the magnitude range $-22.23 \leq M'_r \leq -21.08$. There is no important differences between old and young populations.

Table 4. Median radii in physical units (h^{-1} kpc) from equation (8) for simulated galaxies (see details in Section 5.1), as given by the best-fitting parameters in the M'_r case ($a = 0.75$, $b = -0.36$) and the true total magnitude in the r -band ($a = 0.5$, $b = -0.36$). The results are shown for all objects and, also, split among the old and young populations of simulated galaxies after performing the redefinition of luminosity. The radius $r < 0$ implies $M'_r = M_r$. The ranges in magnitude are those shown in Table 1.

Best-fitting parameters	Ages	$\langle r/h^{-1}\text{kpc} \rangle$	$\langle r/h^{-1}\text{kpc} \rangle$	$\langle r/h^{-1}\text{kpc} \rangle$	$\langle r/h^{-1}\text{kpc} \rangle$
		$M_r = (-19, -18)$	$M_r = (-20, -19)$	$M_r = (-21, -20)$	$M_r = (-22, -21)$
$a = 0.75$, $b = -0.36$	All	716.4	380.6	29.1	< 0
	Old	910.7	723.1	523.4	206.6
	Young	582.2	< 0	< 0	< 0
$a = 0.50$, $b = -0.36$	All	719.9	385.0	28.4	< 0
	Old	902.8	596.7	340.1	6.5
	Young	593.8	36.6	< 0	< 0

all the galaxies, $88 h^{-1}$ kpc for the old galaxies, and $24 h^{-1}$ kpc for the young galaxies. This means that the best-fitting set of parameters that make the assembly bias practically absent for both simulated and real galaxies are truly similar when using luminosity as a proxy of peak height.

The method followed to redefine the luminosity of a galaxy could be compared to adding up the mass from the smooth density field beyond the virial radius. We compare the resulting M'_r luminosity function with that of the total luminosity in haloes in the simulation to help interpret the changes introduced by our approach; this is equivalent to our procedure in Paper I where we compared the mass functions for the original virial mass, and that resulting from our redefinition of peak height, although in this case the luminosity function does not have a theoretical prediction. Figure 18 shows differences in the luminosity function that results from the redefinition of luminosity M'_r for central galaxies (solid circles) and that from adding the luminosity in the r -band of all the galaxies with $M_r < -18$ in the same friends-of-friends (FOF) halo (open triangles) as an approx-

imation of the total luminosity within the peak. The latter is defined for FOF haloes where the magnitude of their central semi-analytic galaxy is $M_r > -22$, otherwise the magnitude of FOF haloes is simply that of their central galaxy. As our formalism adds luminosities of galaxies embedded in haloes, one could have expected the number of these objects with magnitudes M'_r to be similar to that of all the galaxies in FOF haloes, but the difference between the luminosity functions measured with M_r and M'_r is considerable. In this section we will study the origin of this discrepancy.

In Paper I we found that the redefinition based on the mass density field does not change the halo mass function at $M > 10^{12} h^{-1} M_\odot$ (see their Fig. 8) when compared to that of Sheth, Mo, & Tormen (2001, SMT) or that obtained from the virial mass, which is in agreement with the result found by Gao, Springel & White (2005) where the assembly bias with respect to formation time is not important for the high mass regime. The reason for the change found in the low-mass regime is that the peak for an old, low-mass object adds more haloes and mass than for a young object

Table 5. Median radii in physical units (h^{-1} kpc) from equation (8) for SDSS galaxies (see details in Section 5.2), as given by the best-fitting parameters $a = 1$, $b = -0.36$. The results are shown for all objects and, also, split among the old and young populations of observed galaxies after performing the redefinition of luminosity. The radius $r < 0$ implies $M'_r = M_r$. The ranges in magnitude are those shown in Tables 1 and 4.

Best-fitting parameters	Ages	$\langle r/h^{-1}\text{kpc} \rangle$	$\langle r/h^{-1}\text{kpc} \rangle$	$\langle r/h^{-1}\text{kpc} \rangle$	$\langle r/h^{-1}\text{kpc} \rangle$
		$M_r = (-19, -18)$	$M_r = (-20, -19)$	$M_r = (-21, -20)$	$M_r = (-22, -21)$
$a = 1.0, b = -0.36$	All	694.6	358.3	28.2	< 0
	Old	851.8	726.1	535.7	294.5
	Young	606.1	< 0	< 0	< 0




Figure 17. Same as Fig. 12 but using the best-fitting values for the SDSS galaxies $a = 1$ and $b = -0.36$ (solid lines). For comparison, open squares (dashed lines) correspond to the ratio shown in Fig. 11 for the SDSS galaxies using M_r . Note that this comparison is just for reference because the adopted magnitudes are different.

of equal mass. For the following test, instead of considering the mass from dark matter (DM) particles inside a sphere, which corresponds to a continuous field, we use the mass of dark matter haloes in spheres, which is a discrete formalism similar to using the luminosity of individual galaxies. Bear in mind that both estimators measure the neighbours enclosed within the sphere calculating the crowding of the region. In the top and middle panels of Figure 19 we show the correlation function and the velocity infall profile for old and young objects of a given mass for the best-fitting parameters $a = 0.08$ and $b = -0.32$ in Equation (14) of Paper I, resulting from using this discrete approach with neighbour DM haloes. Compared to the best-fit parameters of Paper I ($a = 0.2$ and $b = -0.02$), these are significantly different. The difference between the approach for haloes and that used for a smooth density field is evident when looking at the mass function, shown as open triangles in the bottom panel, which is completely different than that using the




Figure 18. Luminosity function of M'_r magnitudes of semi-analytic galaxies using the parameters $a = 0.75$ and $b = -0.36$ (solid circles). Open triangles correspond to the total r -band luminosity function of FOF haloes given by the contribution from their simulated galaxies with $M_r < -18$.

virial mass (green squares) as well as the predicted mass functions from the extended Press-Schechter model (EPS) and from the SMT model shown as long-dashed and dot-dashed lines, respectively. We also compare it with a set of parameters with good-fitting results to trace the assembly bias using the continuum formalism ($a = 0.2$ and $b = -0.02$, open circles). The behaviour of the latter mass function is in agreement with the virial mass, and the EPS and SMT models.

The discrete formalism actually measures the crowding around objects which constitutes a different proxy of the peak height than the total mass. Figure 20 compares the result obtained with the best-fitting set of values $a = 0.75, b = -0.36$ for the luminosity of individual simulated galaxies and that obtained in Paper I, which considers the mass of DM particles. There is a clear correlation between both methods, showing that the same phenomenon is being described. We calculate a correlation coefficient using the Pearson's product-moment coefficient,




Figure 19. Results using the discrete method for the mass in the simulation. The correlation functions (top) and the velocity profiles (middle) for old and young simulated galaxies ($\delta_t > 0.05$ and $\delta_t < -0.05$, respectively) are similar at large-scales using the best-fitting parameters $a = 0.08$, $b = -0.32$. However, these values do not produce an adequate mass function (triangles, bottom panel) compared with that from the virial mass (squares), the EPS (long-dashed line) and the SMT (dot-dashed line) models, and the set of parameters from Paper I (circles).




Figure 20. Estimates of the overdensity peak height proxy using the approach followed in Paper I that considers the mass of dark matter particles (best-fitting set of parameters $a = 0.2$ and $b = -0.02$, x-axis), and the luminosity of individual simulated galaxies ($a = 0.75$ and $b = -0.36$, y-axis). The mass is in units of the virial mass of the DM substructure that hosts each galaxy. The luminosity is shown as a difference between the original magnitude M_r and M'_r , which is described in Section 5.1. Solid circles are the median and the error bars correspond to the 10 and 90 percentiles of the distribution.

$$C = \frac{\sum_{i=1}^N (\Delta m_i - \overline{\Delta m})(\Delta M - \overline{\Delta M})}{\sqrt{\sum_{i=1}^N (\Delta m_i - \overline{\Delta m})^2 \sum_{i=1}^N (\Delta M - \overline{\Delta M})^2}}, \quad (11)$$

where $\Delta m = M_r - M'_r$ is the difference between the original magnitude M_r and the new one M'_r (y-axis of Fig. 20), $\overline{\Delta m}$ is its average using the N objects, $\Delta M = \log(M/M_{vir})$ is the new mass in units of the original virial mass (x-axis of Fig. 20), and $\overline{\Delta M}$ is its average. We obtain $C = 0.73$, which means that there is a strong correlation between both formalisms.

We conclude that although our method to measure the total luminosity will not reproduce the luminosity function, the reason for this is only the use of discrete neighbours. If we were able to measure the distribution of mass around galaxies and turn it into luminosity, we would obtain a more physical luminosity function, as we did in the case of the mass. This result implies that the phenomena behind the assembly bias, namely, the truncation of mass infall in haloes of low mass embedded in high density regions, can be detected with the new proxy for peak height that can be applied to real galaxy surveys.

7 CONCLUSIONS

We have shown that the assembly bias effect, namely, the difference in the clustering amplitude at large scales of pop-

ulations of equal mass or luminosity but different age, exists for both semi-analytic galaxies and real galaxies from a SDSS catalogue. This is a relevant issue that could affect the ability of the next generation of galaxy surveys to infer accurate cosmological parameters. Using the two-point correlation function, the clustering strength of old SDSS galaxies is from a 25 per cent to a 45 per cent higher than that of young SDSS galaxies of same magnitude at scales $r > 1 h^{-1}$ Mpc. The estimator of age is based on the mass- and luminosity-weighted stellar ages for the synthetic and SDSS galaxies, respectively. In this context, we show that the D_n4000 index is a less adequate parameter to separate active/young and quenched/old galaxies for assembly bias studies using correlation functions at large scales.

Lacerna & Padilla (2011, Paper I) presented an overdensity peak height proxy with the aim to understand the assembly bias effect. This new definition was proposed as a better alternative than the virial mass. In this work we adapt this model including observational issues using galaxy luminosities, taking advantage of their correlation with substructure mass. As the SDSS is limited in flux, and as the tracers of mass are the galaxies themselves, the method is modified to use the crowding of the environment around galaxies in selected volume-limited samples, instead of searching for a correction to the virial mass as was done in Paper I. We measure the total light in cylinders centred in each galaxy, thus obtaining an estimate of the crowding around galaxies, that traces the assembly bias. The radius of this cylinder is parameterized as a function of stellar age and luminosity. The best-fitting sets of parameters in the simulation and observations are very similar ($a = 0.75$, $b = -0.36$ and $a = 1$, $b = -0.36$, respectively). In both cases the assembly bias is practically absent using this approach. The average differences in clustering amplitude between old and young populations for both semi-analytic and observational galaxies are smaller than a 10 percent for each range in magnitude M_r . By estimating a weighted average for the values in the range of 1σ for semi-analytic and SDSS galaxies, it is obtained that the set of best-fitting parameters for the radius of the cylinder is $a = 0.76 \pm 0.31$, $b = -0.39 \pm 0.18$.

This latter model, which constitutes a discrete formalism when counting galaxies within a cylinder or haloes within spheres, does not reproduce the original luminosity or mass functions since it actually measures the crowding around galaxies. However, it is correlated with the smooth density field approach of Paper I. Therefore, both formalisms are associated with the fact that the mass of peaks that did not collapse on to haloes will not be an appropriate proxy for the peak height. It is likely that low-mass haloes in high density environments show their growth truncated by the gravitational effects of their (massive) neighbours. Our results indicate that this also affects the growth of the stellar mass in galaxies in addition to the growth of dark matter haloes. This could induce a misclassification of galaxies in theoretical models to populate haloes, such as HOD, because using virial mass instead of proper peak height.

These results will be of high importance for the next generation of galaxy surveys, such as LSST. The huge amount of information available will allow us to measure the clustering amplitude of galaxies with significant statistics at Mpc scales. Therefore, as we show in this paper, it will be absolutely necessary to model the effect of the environment

on the two-halo regime in order to understand its role in the formation and evolution of haloes and galaxies. Theoretical models will have to include this phenomenon in order to reproduce the observational data with high accuracy in the new era of precision cosmology.

ACKNOWLEDGMENTS

We would like to thank Harold Francke and Alvaro Orsi for comments and discussions. We acknowledge support from FONDAPE “Centro de Astrofísica” 15010003, BASAL-CATA, Fondecyt grant No. 1110328, CONICYT, and MECESUP. Support for IL is provided by the Ministry for the Economy, Development, and Tourism’s Programa Iniciativa Científica Milenio through grant P07-021-F, awarded to The Milky Way Millennium Nucleus. FS thanks supports by the DFG Research Unit FOR 1254. The calculations for this work were performed using the Geryon cluster at the Centre for Astro-Engineering at UC.

Funding for the SDSS and SDSS-II has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the U.S. Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England. The SDSS Web Site is <http://www.sdss.org/>. The SDSS is managed by the Astrophysical Research Consortium for the Participating Institutions. The Participating Institutions are the American Museum of Natural History, Astrophysical Institute Potsdam, University of Basel, University of Cambridge, Case Western Reserve University, University of Chicago, Drexel University, Fermilab, the Institute for Advanced Study, the Japan Participation Group, Johns Hopkins University, the Joint Institute for Nuclear Astrophysics, the Kavli Institute for Particle Astrophysics and Cosmology, the Korean Scientist Group, the Chinese Academy of Sciences (LAMOST), Los Alamos National Laboratory, the Max-Planck-Institute for Astronomy (MPIA), the Max-Planck-Institute for Astrophysics (MPA), New Mexico State University, Ohio State University, University of Pittsburgh, University of Portsmouth, Princeton University, the United States Naval Observatory, and the University of Washington.

The Millennium Simulation databases used in this paper and the web application providing online access to them were constructed as part of the activities of the German Astrophysical Virtual Observatory.

REFERENCES

- Adelman-McCarthy J., et al., 2006, ApJS, 162, 38
- Alonso S., Mesa V., Padilla N., Lambas D. G., 2012, A&A, 539, 46
- Balogh M., Morris S., Yee H. K. C., Carlberg R., Ellingson E., 1999, ApJ, 527, 54
- Blanton M. R., Roweis S., 2007, AJ, 133, 734
- Colless M., et al., 2001, MNRAS, 328, 1039
- Conroy C., Wechsler R., Kravtsov A., 2006, ApJ, 647, 201
- Cooper M. C., Gallazzi A., Newman J. A., Yan R., 2010, MNRAS, 402, 1942

- Croton D. J., Gao L., White S. D. M., 2007, MNRAS, 374, 1303
- Dalal N., White M., Bond J. R., Shirokov A., 2008, ApJ, 687, 12
- Davis M., Efstathiou G., Frenk C., White S. D. M., 1985, ApJ, 292, 371
- De Lucia G., Blaizot J., 2007, MNRAS, 375, 2
- Faltenbacher A., White S. D. M., 2010, ApJ, 708, 469
- Fukugita M., Ichikawa T., Gunn J. E., Doi M., Shimakasu K., Schneider D. P., 1996, AJ, 111, 1748
- Gallazzi A., Charlot S., Brinchmann J., White S. D. M., Tremonti C. A., 2005, MNRAS, 362, 41
- Gao L., Springel V., White S. D. M., 2005, MNRAS, 363, 66
- Gao L., White S. D. M., 2007, MNRAS, 377, L5
- Guo Q., et al., 2011, MNRAS, 413, 101
- Hahn O., Porciani C., Dekel A., Carollo C. M., 2009, MNRAS, 398, 1742
- Jester S., et al., 2005, AJ, 130, 873
- Komatsu E., et al., 2011, ApJS, 192, 18
- Lacerna I., Padilla N., 2011, MNRAS, 412, 1283 (Paper I)
- Lagos C., Cora S. A., Padilla N., 2008, MNRAS, 388, 587
- Lagos C., Padilla N., Cora S., 2009, MNRAS, 395, 625
- LSST Science Collaborations, Abell P., et al., 2009, arXiv:0912.0201
- Muldrew S., et al., 2012, MNRAS, 419, 2670
- Paz D. J., Stasyszyn F., Padilla N., 2008, MNRAS, 389, 1127
- Saunders, W., Rowan-Robinson, M., Lawrence, A., 1992, MNRAS, 258, 134
- Sheth R. K., Mo H. J., Tormen G., 2001, MNRAS, 323, 1
- Simha V., Weinberg D. H., Davé R., Fardal M., Katz N., Oppenheimer B. D., 2012, MNRAS, 423, 3458
- Skibba R., Sheth R. K., Connolly A. J., Scranton R., 2006, MNRAS, 369, 68
- Smith J. A., et al., 2002, AJ, 123, 2121
- Springel V., et al., 2005, Nature, 435, 629
- Springel V., White S. D. M., Tormen G., Kauffmann G., 2001, MNRAS, 328, 726
- Tinker J. L., Wetzel A. R., Conroy C., 2011, arXiv:1107.5046
- Wang H., Mo H. J., Jing Y. P., 2007, MNRAS, 375, 633
- Wang Y., Yang X., Mo H. J., van den Bosch F., Weinmann S., Chu Y., 2008, ApJ, 687, 919
- Wechsler R. H., Zentner A. R., Bullock J. S., Kravtsov A. V., Allgood B., 2006, ApJ, 652, 71
- Wu H., Rozo E., Wechsler R. H., 2008, ApJ, 688, 729
- Yang X., Mo H. J., van den Bosch F. C., 2006, ApJ, 638, L55
- York D. G., et al., 2000, AJ, 120, 1579
- Zapata T., Perez J., Padilla N., Tissera P., 2009, MNRAS, 394, 2229
- Zehavi I., et al., 2005, ApJ, 630, 1
- Zhu G., et al., 2006, ApJ, 639, L5